

(to appear in the Astronomical Journal)

The Universality of the Fundamental Plane of E and S0 Galaxies. Spectroscopic Data¹

Marco Scodeggio², Riccardo Giovanelli and Martha P. Haynes

Center for Radiophysics and Space Research and National Astronomy and Ionosphere Center³,
Cornell University, Ithaca, NY 14853

mcodegg@eso.org

riccardo, haynes@astrosun.tn.cornell.edu

ABSTRACT

We present here central velocity dispersion measurements for 325 early-type galaxies in eight clusters and groups of galaxies, including new observations for 212 galaxies. The clusters and groups are the A262, A1367, Coma (A1656), A2634, Cancer and Pegasus clusters, and the NGC 383 and NGC 507 groups. The new measurements were derived from medium dispersion spectra, that cover 600 Å centered on the Mg Ib triplet at $\lambda \sim 5175$. Velocity dispersions were measured using the Tonry & Davis cross-correlation method, with a typical accuracy of 6%. A detailed comparison with other data sources is made.

Subject headings: distance scale — galaxies: distances and redshifts — galaxies: elliptical and lenticular — galaxies: fundamental parameters — galaxies: kinematics and dynamics

1. Introduction

This is the fourth in a series of papers (Scodeggio, Giovanelli & Haynes 1997a,b; hereafter Paper I and Paper II; Scodeggio, Giovanelli & Haynes 1998a; hereafter Paper III) devoted to a

¹ Work based in part on observations obtained with the Hale 200" telescope at the Palomar Observatory. These observations were made as part of a continuing collaborative agreement between the California Institute of Technology and Cornell University.

²Present address: European Southern Observatory, Karl-Schwarzschild-Straße 2, D-85748, Garching bei München, Germany.

³The National Astronomy and Ionosphere Center is operated by Cornell University under a cooperative agreement with the National Science Foundation.

study of the universality of the Fundamental Plane (FP) relation of early-type galaxies (Djorgovski & Davis 1987; Dressler *et al.* 1987). The universality of the FP, and of other distance indicator relations like the Tully-Fisher (TF) relation (Tully & Fisher 1977), is the fundamental assumption behind their use, but this assumption cannot be taken for granted, because of the existence of well known environmental effects that can link the evolution of a galaxy to the properties of its surrounding environment. It is therefore important to obtain an observational verification of such universality (discussions on this subject are presented by Djorgovski, de Carvalho & Han 1988; Guzmán *et al.* 1992; Jørgensen, Franx & Kjaergaard 1996; Schroeder 1996; Giovanelli *et al.* 1997b; Paper I).

In this series, we present the results of a FP study of eight clusters, with the aim of obtaining a quantitative verification of the FP universality through a comparison of accurate distance estimates obtained using both the FP and the TF relations. These clusters are taken from the list of clusters studied by Giovanelli *et al.* (1997a,b) using the TF relation, and are the Coma, A1367, A2634, A262, Cancer, and Pegasus clusters, and the NGC 507 and NGC 383 groups. In this paper we present spectroscopic data, necessary for building the FP relation, for 325 galaxies, including new observations for 212 galaxies, while Paper III contains I band CCD photometric data, and describes the cluster selection and membership criteria adopted for the FP work. In successive work (Scodreggio, Giovanelli & Haynes 1998b) we will discuss the properties of the I band FP relation, and the comparison between TF and FP distance estimates for the clusters in our sample.

This paper is organized as follows: sections 2, 3, and 4 present our new spectroscopic observations and the adopted data reduction procedures. Sections 5 and 6 present the complete spectroscopic sample that will be used in the FP work, and the internal and external comparisons used to assess the quality of our new measurements.

2. Observations

New spectroscopic observations for this work were obtained during 8 observing runs with the Hale 5m telescope of the Palomar Observatory, from September 1992 to September 1996. Table 1 summarizes these observing runs, and gives the parameters of the relative instrumental setups. During all runs the red camera of the Double Spectrograph (Oke & Gunn 1982) was used with a 1200 lines/mm grating, to produce spectra with a dispersion of approximately 0.82 Å per pixel (except in runs 7 and 8, when a different CCD chip was mounted on the camera, and the dispersion was approximately 0.66 Å per pixel). This resulted in a spectral coverage of approximately 600 Å, from 5000 to 5600 Å, including the Mg Ib lines at $\lambda \sim 5175$. The spectral resolution, measured both from the width of the calibration lamps lines, and from the width of the peak in the autocorrelation function of the spectrum of velocity standard stars, was approximately 2.2 Å (FWHM) for all runs. This is equivalent to a velocity resolution of 129 km s⁻¹ at 5300 Å. All observations were obtained with a 2 arcsec wide slit, while the length of the slit was 128 arcsec. The slit orientation was kept in the East-West direction, unless a close pair of galaxies

could be observed at once with the proper slit orientation. Galaxies were visually centered in the slit using the acquisition camera of the Double Spectrograph. Integration times varied between 10 and 90 minutes, depending on the brightness of the galaxy. The median integration time was 30 minutes. He-Ne-Ar lamp spectra were obtained before and after each galaxy observation, to provide wavelength calibration. Late G and early K type giant stars were observed each night to serve as templates for radial velocity and velocity dispersion determinations. To maximize the signal-to-noise ratio in these template spectra, the star observations were obtained with 30-120 sec integrations, while trailing the star along the slit. This allowed to spread the light on a large area of the CCD chip without saturating the image.

3. Basic data reduction and wavelength calibration

The data reduction process was performed entirely using standard IRAF⁴ procedures. All frames were bias-subtracted and flat-fielded, using dome flats. The number of pixels affected by cosmic rays hits was small in all the frames, and they were blanked during a phase of visual inspection of all frames. Because most of the spectroscopic observations were obtained in non photometric conditions, no attempt was made to obtain a flux calibration for any of the derived spectra.

Wavelength calibration was performed in a two-step process. First, the He-Ne-Ar lamp spectra frames obtained before and after each galaxy spectrum were used to obtain a two-dimensional dispersion solution, where a wavelength value is assigned to each pixel in the CCD frame. Typically 12-14 spectral lines, spread uniformly over the entire wavelength range, were used in the fit. The rms deviations between fitted and true wavelength were of the order of 0.06–0.1 Å in the central portion of the frames (where the galaxy spectra are positioned), and increased up to 0.25 Å at the edges of the frame, due to the camera astigmatism and to focusing problems caused by the less than perfect flatness of the CCD chip. Single line deviations from the best fit solutions were always less than 0.3 Å. An uncertainty in the wavelength calibration of 0.1 Å at 5300 Å translates into a velocity uncertainty of 5.6 km s^{−1}. The CCD chip used for runs 7 and 8 does not have focusing problems, and therefore the rms uncertainty in the calibration of the spectra obtained in those runs was of the order of 0.08 Å, uniformly across the entire frame. The two-dimensional dispersion solutions were applied to the galaxy spectral frames, so that all pixels in one row of the corrected CCD frame were characterized by the same wavelength. Finally the two strong night-sky lines at 5460.7 Å (Hg) and 5577.4 Å (OI) were used to check the absolute wavelength calibration. The 1-σ dispersion in the measured wavelengths of these lines was approximately 0.08 Å. A small fraction (2-3%) of the frames showed large deviations ($> 4\sigma$) in the wavelength measured for the

⁴IRAF (Image Reduction and Analysis Facility) is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy (AURA), Inc., under a cooperative agreement with the National Science Foundation.

two sky lines. For these cases the original dispersion solution was rigidly shifted to bring the two lines into agreement with their expected wavelengths.

An independent check on the wavelength calibration was performed using the velocity standard star spectra. Radial velocities were measured repeatedly using all star spectra obtained in the different runs. Using in turn one star as velocity standard and all others as unknown, radial velocities were derived, to be compared with the known radial velocity of each star. The rms deviations between the measured and the known velocities were between 6.0 and 8.5 km s^{-1} , when spectra obtained during a single run only were considered, and approximately 9 km s^{-1} when spectra from all runs were considered. This result is in good agreement with the calibration uncertainty derived from the sky lines measurements. Therefore 9 km s^{-1} is an estimate of the overall accuracy we can achieve with this spectroscopic data set.

4. Determination of the spectroscopic parameters

During the last twenty years a number of methods have been developed to perform velocity dispersion measurements as objectively as possible (see for example Sargent *et al.* 1977; Tonry & Davis 1979; Franx, Illingworth & Heckman 1989; Bender 1990). These methods are all based on the comparison between the spectrum of the galaxy whose velocity dispersion is to be determined, and a fiducial spectral template. This can be either the spectrum of an appropriate star, with spectral lines unresolved at the spectral resolution being used, or a high signal-to-noise spectrum of a galaxy with known velocity dispersion. There are small but in principle significant differences among these methods, but in practice all methods seem to give results in good relative agreement, when applied to spectra with high signal-to-noise ratio, making the choice of one particular method a non critical issue.

In this work, quantitative measurements on the spectra have been obtained using the cross-correlation technique of Tonry & Davis (1979), implemented in the IRAF task *fxcor*. The basic assumption behind this and other similar methods is that the spectrum of an elliptical galaxy (and also of the bulge of a disk galaxy) is well approximated by the spectrum of its most luminous stars (K0–K1 giants), modified only by the effects of the stellar motions inside the galaxy. Since these motions introduce just a Doppler shift in the stellar spectra, the galaxy spectrum is given by the convolution of the spectrum of a K giant star with the line of sight stellar velocity distribution (LOSVD). Therefore the LOSVD can be obtained with a deconvolution process from the galaxy spectrum and a suitable stellar template. Since cross-correlation and convolution are two related operators, the Tonry & Davis algorithm uses cross-correlation as a computational tool to derive the LOSVD. The procedure takes advantage of the fact that the Fourier transform of a convolution, or of a cross-correlation, reduces to a product between individual functions’ transforms.

The one-dimensional galaxy spectra used for the radial velocity and velocity dispersion measurements were extracted from the two-dimensional wavelength calibrated frames using a 6

arcsec wide window, centered on the peak of the galaxy continuum light. This, combined with the 2 arcsec slit aperture, provides a final effective aperture of 6×2 arcsec for all the galaxy observations. The stellar templates were obtained averaging the signal over 20–30 pixels in the spatial dimension at all wavelengths, to produce spectra with a signal to noise ratio greater than 200. Before computing the Fourier transforms and the cross-correlation function, all spectra were continuum-subtracted, normalized, end-masked with a cosine bell function, and re-binned to a logarithmic wavelength scale. The final velocity dispersion measurements have been obtained by averaging the determinations obtained with five different stellar templates, of spectral types between G9 and K2. No significant dependence of the measurements on the template star spectral type was observed.

4.1. The Effect of the Noise

Because of the limited signal-to-noise (S/N) ratio of the spectra being analyzed, and also to facilitate comparisons with other studies, we have constrained the deconvolution procedure by assuming the LOSVD to be Gaussian, and therefore characterized by two parameters only: a mean redshift z and a velocity dispersion σ . The reliability of this method has been tested using simulations with synthetic spectra. Stellar templates have been broadened with Gaussian profiles of known width to simulate a large range of velocity dispersions, and noise has been added to the resulting spectra, to reproduce the S/N characteristic of the galaxy spectra. Then velocity dispersions have been measured using the original stellar spectrum as a template. These tests show that *fxcor* produces an overestimate of the velocity dispersion, at the 4-5% level, while no bias is present in the measurement of radial velocities. The raw measurements have been therefore corrected to remove this effect. To a first approximation, this effect appears to be independent of the S/N ratio of the input spectrum, but no simulation was performed using spectra with S/N ratio lower than 20.

If the true LOSVD is approximately Gaussian, then the statistical uncertainty with which the velocity dispersion is obtained is a function of the S/N ratio of the galaxy spectrum only, since the S/N ratio in the template spectrum is many times higher than that of the galaxy spectrum (see Tonry & Davis 1979 for details). An estimate of the amplitude of this uncertainty was obtained using synthetic spectra with added noise. It was found that a good approximation to the value of the uncertainty is given by the relation $\epsilon_{\log \sigma} \simeq 1.1/(S/N)$. The median uncertainty on the radial velocity determination is 13 km s^{-1} , and on the velocity dispersion is 6%.

As a visual guide to the quality of the data presented in this work, we show in Figure 1 the spectra of 6 different galaxies. These are divided into two groups, and all three galaxies in one group have very similar velocity dispersion (approximately 140 and 230 km s^{-1} , respectively), but their spectra have different S/N ratio. These spectra were chosen to provide an example of a spectrum at the upper quartile, median, and lower quartile in the distribution of S/N values for the spectra in our dataset.

4.2. The Effect of Rotation

Galaxy one-dimensional spectra were extracted using a 6 arcsec aperture, which can bracket a fairly large portion of the target galaxy, especially for more distant clusters. Therefore galaxy rotation, if present, can contribute some broadening of the spectral lines and bias the velocity dispersion measurements. We thus use the galaxy two-dimensional spectra to estimate the galaxy rotation velocity, and gauge whether a correction of the velocity dispersion measurements is necessary. A 5 point rotation curve within the 6 arcsec aperture (10 pixels) has been obtained for each galaxy, using 5 one-dimensional spectra, extracted using 2-pixel wide apertures positioned side by side. Using the cross-correlation method, individual mean velocities can be obtained for the 5 spectra. The differences in the radial velocities measured between each one of the four lateral spectra and the one measured in the central 2 pixels define the projection of the galaxy rotation curve along the slit.

The contribution of a given rotation to the broadening of the spectral lines has been estimated using a very simple model. A galaxy spectrum has been simulated combining 5 copies of a stellar template, broadened with a Gaussian of fixed width, and shifted to reproduce the 5 velocities in the rotation curve. The weights used in the combination have been derived from the relative intensities in the continuum of the 5 one-dimensional galaxy spectra used to obtain the rotation curve. The comparison of the velocity dispersion in the simulated spectrum with the one used to broaden the stellar template spectra provides an estimate of the amount of broadening due to the rotation.

Given the accuracy with which radial velocities can be measured, and the small number of points used to define the rotation curve, rotation velocities smaller than $\simeq 20 \text{ km s}^{-1}$ cannot be reliably measured. Within the sample presented here, 75 galaxies do not show detectable rotation. The remaining 137 have a median rotation velocity of 40 km s^{-1} , with the largest measured velocity being 130 km s^{-1} . Note that we had no a priori information on the position angle of the axis of rotation of each galaxy. This rotation is responsible for an average broadening of the LOSVD of $\simeq 4\%$, with a maximum of $\simeq 9\%$ for the largest rotation velocities.

4.3. Aperture Correction

Because of radial gradients in the stellar velocity dispersion within early-type galaxies, the velocity dispersion measured in a given galaxy depends on the size of the aperture that is used to extract its spectrum. When spectra are obtained with a fixed aperture size (in arcsec), this aperture covers regions of different physical size (in kiloparsec) on galaxies at different distances, and this produces biased velocity dispersion measurements. Therefore fixed aperture measurements must be corrected to a standard, distance independent, aperture size. Here all measurements are corrected to the value one would obtain for a galaxy at the distance of the

Coma cluster, using the relation derived by Jørgensen, Franx & Kjaergaard (1995)

$$\log \frac{\sigma_{ap}}{\sigma_{norm}} = -0.04 \log \frac{r_{ap}}{r_{norm}} = -0.04 \log \frac{d}{d_{Coma}} \quad (1)$$

where σ_{ap} and σ_{norm} are the measured and corrected velocity dispersion, r_{ap} and r_{norm} are the projected angular size of a given length at the distance of the observed galaxy and at the reference distance used for the normalization, and d and d_{Coma} are the distance of the cluster to which the given galaxy belongs, and that of the Coma cluster. Distances are derived from the cluster redshifts, in the CMB reference frame.

5. The spectroscopic sample

Spectroscopic data for early-type galaxies in 8 clusters and rich groups of galaxies are presented here. These are the NGC 383 group, NGC 507 group, A262, Cancer, A1367, Coma (A1656), Pegasus, A2634. A short discussion on their properties, and on the membership criteria we used to assign galaxies to these clusters, is presented in Paper III. Our spectroscopic sample is composed of two partly overlapping datasets: our own observations, discussed in the previous sections, for a total of 212 galaxies, and a compilation of data from the literature, which includes 152 galaxies. For the Coma cluster, most of the data come from the literature, while for the other clusters most of the data are new. In all cases we have tried to complement the existing observations in order to obtain approximately flux limited samples, while having enough overlap with those observations, as to compare the different velocity dispersion scales.

5.1. New observations

The criteria adopted to select galaxies for our spectroscopic observations were as follows: CGCG magnitudes (Zwicky *et al.* 1963-1968) were used for the sample selection, if available. Otherwise our own eye estimates, calibrated on the same scale, were used. All galaxies with morphological type E, S0, and S0a were included in the sample. A few objects, that were originally classified as S0 or S0a, were later classified Sa on the basis of the CCD imaging (see Paper III), but they were left in the sample if a spectrum had already been taken. Only galaxies with previously available redshift measurements, so that their cluster membership was already known, were targeted for observations.

No particular criterion was used to select the objects that would provide an overlap with already available datasets. An effort was made, however, to cover as large an interval of velocity dispersions as possible.

Table 2 presents the relevant parameters for the 212 objects for which we have obtained spectroscopic observations. Only the first page is presented herefor guidance regarding its form

and content . The table is available in its entirety in digital form on the Journal Electronic Edition, or from the authors. The table is organized as follows:

Col. 1: Galaxy name: if the galaxy is listed in the UGC catalog (Nilson 1973), the UGC number is given; if not, our internal coding number, from the private galaxy catalog of Giovanelli & Haynes referred to as the AGC, is given.

Cols. 2 and 3: the NGC or IC number, if available, and the CGCG field and ordinal number within that field, if the galaxy is listed in the CGCG (Zwicky *et al.* 1963-1968). For galaxies in the Coma cluster that are not listed in either one of those catalogs, the Dressler number (Dressler 1980) is given, if available.

Cols. 4 and 5: Right Ascension and Declination, in 1950.0 epoch.

Col. 6: morphological type code, in the RC3 (de Vaucouleurs *et al.* 1991) scheme (-5 for E, -2 for S0, 0 for S0a). Morphological types were derived from joint inspection of the blue plates of the Palomar Observatory Sky Survey (POSS), and of the CCD images discussed in Paper III.

Col. 7: Measured heliocentric velocity, in km s^{-1} .

Col. 8: Measured stellar velocity dispersion, in km s^{-1} .

Col. 9: Measured rotation velocity, in km s^{-1} . If no data is present, the spectrum had to low a S/N ratio to perform the measurement.

Col. 10: Velocity dispersion, in km s^{-1} , corrected for galaxy rotation.

Col. 11: Logarithm of the rotation-corrected velocity dispersion.

Col. 12: Uncertainty on the logarithm of the velocity dispersion.

Col. 13: Aperture-corrected logarithm of the velocity dispersion.

Col. 14: A reference code indicates if other velocity dispersion measurements were available in the literature, and their source. If an asterisk appears in this column, special comments on the object are included at the foot of the table.

A subset of these data has already been published in Paper I, limited to galaxies in the Coma and A2634 clusters. Since then, a few galaxies have been reobserved, and the data combined to obtain new measurements of the relevant photometric parameters. The data presented here therefore supersede that previous report.

5.2. Data from the literature

Velocity dispersion measurements are available in the literature for each of the clusters in our sample. The number is large only in the case of the Coma cluster, where measurements from Dressler (1987), Davies *et al.* (1987), Faber *et al.* (1989), and Lucey *et al.* (1991) are available for a total of 103 galaxies, and in the case of A2634, where measurements from Lucey *et al.* (1997) are available for 37 galaxies. For the remaining clusters, we have found data in Faber *et al.* (1989) for 6 galaxies in A262, 7 galaxies in A1367, 1 galaxy in Cancer, 4 galaxies in Pegasus, 12 galaxies in the NGC 507 and NGC 383 groups. For the brightest galaxy in some of the clusters multiple measurements from various sources are available (see the footnotes to Table 2). In all clusters, we

have limited our search to galaxies for which we have obtained I band photometric measurements (see Paper III). Therefore a larger spectroscopic dataset than the one reported here might be available in some cases. Table 3 lists the relevant parameters for the 170 objects for which data are derived from the literature. Only the first page is presented here; the entire table is available in digital form on the Journal Electronic Edition, or from the authors. The table is organized as follows:

Cols. 1 to 6: same as in Table 2.

Col. 7: Heliocentric velocity, in km s^{-1} .

Col. 8: Stellar velocity dispersion, in km s^{-1} .

Col. 9: Logarithm of the velocity dispersion.

Col. 10: Uncertainty on the logarithm of the velocity dispersion. A common value for each main dataset was adopted, corresponding to the typical uncertainty quoted by the authors.

Col. 11: Aperture-corrected logarithm of the velocity dispersion.

Col. 12: Reference code.

6. Internal and external comparison

Repeated observations of a fraction of the galaxies in our sample provide the opportunity to evaluate the internal consistency of the spectroscopic measurements. However, the number of repeated observations is rather small, because spectroscopic observations are quite expensive in terms of telescope time. In total, 27 pairs of observations are available for an internal comparison. Table 4 lists the results of the comparison of radial velocity and velocity dispersion. Mean statistical uncertainties associated with the derivation of those parameters are included in the Table, multiplied by a $\sqrt{2}$ factor to allow a direct comparison with the rms scatter in the observations (assuming equal error contributions to the rms scatter). The rms scatter in the radial velocity comparison (see Figure 2a) is surprisingly large (29 km s^{-1}), but it is mostly due to just 2 galaxies. After removing these extreme cases the scatter is reduced to 19 km s^{-1} , in agreement with the statistical uncertainties on the radial velocity determinations. Unfortunately with such small number statistics it is difficult to decide whether these 2 objects are truly outliers. However we point out that in both cases one of the two spectra being compared has an extremely low signal-to-noise ratio, and therefore the uncertainty on the determination of the corresponding spectroscopic parameters is expected to be large.

The comparison of the velocity dispersions is presented in Figure 2b, and shows that, as expected, the accuracy of the dispersion determinations for very low velocity dispersion objects ($\sigma < 100 \text{ km s}^{-1}$) is lower than that for higher velocity dispersion objects. The rms scatter measured for the 6 low velocity dispersion objects is in fact more than twice as large as the one measured for the remaining 21 objects. When these low velocity dispersion objects are removed from the sample, we find a fair agreement between the expected uncertainties and the observed dispersion in the measurements.

The absolute consistency of the velocity dispersion measurements is evaluated through a comparison with measurements taken from the literature. There are 62 observations for 57 galaxies for which this comparison is possible. The results of such a comparison are presented in Table 5 and in Figure 3, where the difference (us - literature) is plotted vs. the mean value of the velocity dispersion, derived combining the two measurements. Five extremely discrepant objects are removed from this comparison. For two objects the measurement we have obtained for their velocity dispersion is affected by a large uncertainty. For the remaining three no easy explanation for the observed discrepancy is available. It is clear that there is good agreement with the observations of Dressler (1987), Davies *et al.* (1987), and Faber *et al.* (1989), collectively indicated as 7S in Table 5, and also with the observations of Lucey *et al.* (1997). No significant offset is present, and the observed rms scatter of the differences in the values of $\log \sigma$ is in good agreement with an expected one, based on the 6% median uncertainty of our measurements, and on an average 10% uncertainty for the measurements in the literature. A small discrepancy is present when the Coma cluster observations of Lucey *et al.* (1991) are considered, but the sample is probably too small to derive robust conclusions. We notice that the comparison between Lucey *et al.* (1991) and Faber *et al.* (1989) datasets, based on a larger sample of galaxies, shows good relative agreement (see Lucey *et al.* 1991). Given the good agreement between the different datasets, all measurements presented in Table 2 and 3 will be merged for the FP work that will be discussed in Scodeggio *et al.* (1998b).

We would like to thank John Henning, Mike Doyle, Dave Tennant, and Bob Thicksten for their technical support during the various observing runs at Palomar, Juan Carrasco for his masterful handling of many observing nights, and Gary Wegner for helpful discussions on the comparison between different spectroscopic datasets. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research is part of the Ph.D. Thesis of MS, and is supported by the NSF grants AST94–20505 and AST96–17069 to RG and AST95–28860 to MPH.

REFERENCES

- Bender, R. 1990, A&A, 229, 441
- Davies, R.L., & Illingworth, G. 1983, ApJ, 266, 516
- Davies, R.L., Burstein, D., Dressler, A., Faber, S.M., Lynden-Bell, D., Terlevich, R.J., & Wegner, G. 1987, ApJS, 64, 581
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H.G., Buta, R., Paturel, G., & Fouqué, P. 1991, “Third Reference Catalogue of Bright Galaxies” (New York: Springer) (RC3)
- Djorgovski, S.G., & Davis, M. 1987, ApJ, 313, 59

- Djorgovski, S.G., De Carvalho, R.R., & Han, M.S. 1988, in “The Extragalactic Distance Scale”, ed. S. van den Bergh & J. Pritchet, p. 329
- Dressler, A. 1980, *ApJS*, 42, 565
- Dressler, A. 1987, *ApJ*, 317, 1
- Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R.L., Faber, S.M., Terlevich, R.J., & Wegner, G. 1987, *ApJ*, 313, 42
- Faber, S. M., & Jackson, R. E. 1976, *ApJ*, 204, 668
- Faber, S.M., Wegner, G., Burstein, D., Davies, R.L., Dressler, A., Lynden-Bell, D., & Terlevich, R.J. 1989, *ApJS*, 71, 173
- Fisher, D., Illingworth, G., & Franx, M. 1995, *ApJ*, 438, 539
- Franx, M., Illingworth, G., & Heckman, T. 1989, *ApJ*, 344, 613
- Giovanelli, R., Haynes, M.P., Herter, T., Vogt, N.P., da Costa, L.N., Freudling, W., Salzer, J.J. and Wegner, G. 1997a, *AJ*, 113, 22 (G97a)
- Giovanelli, R., Haynes, M.P., Herter, T., Vogt, N.P., da Costa, L.N., Freudling, W., Salzer, J.J. and Wegner, G. 1997b, *AJ*, 113, 53 (G97b)
- Guzmán, R., Lucey, J.R., Carter, D., & Terlevich, R.J. 1992, *MNRAS*, 257, 187
- Jørgensen, I., Franx, M., & Kjaergaard, P. 1995, *MNRAS*, 276, 1341
- Jørgensen, I., Franx, M., & Kjaergaard, P. 1996, *MNRAS*, 280, 167
- Lucey, J.R., Guzmán, R., Carter, D., & Terlevich, R.J. 1991, *MNRAS*, 253, 584
- Lucey, J.R., Guzmán, R., Steel, J., & Carter, D., 1997, *MNRAS*, 287, 899
- Malumuth, E.M., & Kirshner, R.P. 1981, *ApJ*, 251, 508
- Nilson, P. 1973, “Uppsala General Catalogue of Galaxies” (Uppsala Astron. Obs. Ann., Vol. 6) (UGC)
- Oegerle, W.R., & Hoessel, J.G. 1991, *ApJ*, 375, 15
- Oke, J.B., & Gunn, J.E. 1982, *PASP*, 94, 586
- Sargent, W.L.W., Schechter, P.L., Boksenberg, A., & Shortridge, K., 1977, *ApJ*, 212, 326
- Schroeder, A., 1996, PhD Thesis, Univ. of Basel
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1997a, *AJ*, 113, 101 (Paper I)
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1997b, *AJ*, 113, 2087 (Paper II)
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1998a, submitted to *AJ* (Paper III)
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1998b, in preparation
- Tonry, J.L. 1984, *ApJ*, 279, 13
- Tonry, J.L. 1985, *AJ*, 90, 2431

Tonry, J.L., & Davis, M. 1979, AJ, 84, 1511

Tonry, J.L., & Davis, M. 1981, ApJ, 246, 666

Tully, R.B., & Fisher, J.R. 1977, A&A, 54, 661

Zwicky, F., Herzog, E., Wild, P., Karpowicz, M., & Kowal, C. 1961–1968, “Catalogue of Galaxies and Clusters of Galaxies” (Pasadena: California Institute of Technology)

Table 1: Spectroscopic observing runs

	Runs 1–6	Runs 7–8
Telescope	Palomar 5m	Palomar 5m
Instrument	DS-red camera	DS-red camera
CCD	TI 305 #8	Tek1024
Pixels	800 x 300	1024 x 300
Read-out noise	11 e [−]	7.5 e [−]
Gain	1.7 e [−] /DU	2.0 e [−] /DU
Spatial scale	0.61 ″/pixel	0.46 ″/pixel
Slit aperture	2 arcsec	2 arcsec
Grating	1200 l/mm	1200 l/mm
Dispersion	0.82 Å/pixel	0.66 Å/pixel
Spectr. resol.	2.2 Å (129 km/s)	2.2 Å (129 km/s)
Wavelength cover.	4960-5620 Å	4960-5660 Å

	Run 1	Run 2	Run 3	Run 4
Dates	Sep. 23-26 1992	Sep. 17-20 1993	Mar. 3-6 1994	Apr. 28-29 1994
Nights used	2	4	3	1
Nr. of galaxies	16	64	31	6

	Run 5	Run 6	Run 7	Run 8
Dates	Sep. 6-12 1994	Aug. 23-25 1995	Mar. 19-20 1996	Sep. 20 1996
Nights used	7	3	1	1
Nr. of galaxies	72	28	5	8

Table 4. Spectroscopic data quality: internal comparison

Parameter	N_{gal}	mean difference	rms scatter	mean uncertainty
V_{hel} (all)	27	5.8	28.9	18.0
V_{hel} (2 most discr. excluded)	25	-2.2	19.3	18.0
$\log \sigma$ (all)	27	0.002	0.055	0.031
$\log \sigma$ ($\sigma > 100 \text{ km s}^{-1}$)	21	0.002	0.041	0.037
$\log \sigma$ ($\sigma < 100 \text{ km s}^{-1}$)	6	0.001	0.096	0.037

Table 5. Spectroscopic data quality: external comparison

Parameter	N_{gal}	mean difference	rms scatter
$\log \sigma$ (all)	56	-0.004	0.053
$\log \sigma$ (7S)	29	-0.004	0.050
$\log \sigma$ (L97)	21	-0.001	0.057
$\log \sigma$ (L91)	6	-0.015	0.056




Fig. 1.— Spectra of 6 different galaxies in our data-set. These are divided into two groups, and all three galaxies in one group have very similar velocity dispersion (approximately 140 and 230 km s^{-1} , respectively), but their spectra have different S/N ratio. These spectra were chosen to provide an example of a spectrum at the upper quartile, median, and lower quartile (top to bottom) in the distribution of S/N values for the spectra in our data-set.




Fig. 1.— (Continued)




Fig. 2.— Internal comparison of the newly derived spectroscopic parameters. ΔV_{hel} and $\Delta \log_{10} \sigma$ are the differences between the values of the heliocentric radial velocity and of the logarithm of the velocity dispersion obtained from repeated observations of 27 galaxies. They are plotted as a function of the mean value of the relative parameter, obtained combining the different measurements.




Fig. 3.— External comparison of the velocity dispersion measurements for galaxies in our sample. $\Delta \log_{10} \sigma$ is the difference between the value of the logarithm of the velocity dispersion we obtained for a given galaxy, and the value found in the literature. This difference is plotted against mean value of this parameter, obtained combining the different measurements. Filled circles identify the comparison with the measurements of Faber *et al.* (1989), filled triangles the comparison with the measurements of Lucey *et al.* (1997), while open squares identify the comparison with the data of Lucey *et al.* (1991).

TABLE 2. Measred spectroscopic parameters

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
U680	N374	501-080	1 04 19.6	32 31 46	0	4991	177	0	177	2.248	0.019	2.241	
A110036	N375	...	1 04 20.2	32 04 52	-5	5901	175	102	161	2.207	0.018	2.200	
U683	N379	501-082	1 04 30.0	32 15 16	-2	5486	228	20	227	2.356	0.022	2.349	1
U682	N380	501-081	1 04 32.3	32 13 01	-5	4426	325	141	310	2.491	0.016	2.485	1
U688	N382	501-086	1 04 38.7	32 08 13	-5	5220	198	31	197	2.294	0.020	2.288	1
U689	N383	501-087	1 04 39.4	32 08 46	-2	5056	272	55	270	2.431	0.017	2.425	1
U686	N384	501-084	1 04 39.7	32 01 33	-2	4251	254	95	246	2.391	0.018	2.384	
U687	N385	501-085	1 04 41.9	32 03 15	-5	4994	203	18	202	2.305	0.019	2.299	1
A110044	N386	501-088	1 04 45.5	32 05 43	-5	5517	120	0	120	2.079	0.028	2.072	1
A110047	N388	501-090	1 05 01.4	32 02 36	-2	5431	135	0	135	2.130	0.028	2.124	
U700	N392	501-094	1 05 37.1	32 52 00	-2	4674	245	62	242	2.384	0.015	2.377	1
A100055	N394	501-095	1 05 39.7	32 52 53	-2	4390	177	0	177	2.248	0.019	2.241	
A110057	N397	501-096	1 05 44.8	32 50 34	-2	4966	146	38	143	2.155	0.021	2.149	
U712	N399	501-101	1 06 13.0	32 22 06	0	5231	131	20	130	2.114	0.020	2.107	
U715	N403	501-104	1 06 28.2	32 29 05	0	5120	215	108	204	2.310	0.018	2.303	
U730	N407	501-115	1 07 49.9	32 51 32	-5	5564	221	0	221	2.344	0.024	2.338	
U735	N410	501-118	1 08 11.6	32 53 10	-5	5274	287	18	287	2.458	0.014	2.451	1
A110098	...	501-119	1 08 20.8	32 57 41	-5	4742	143	58	138	2.140	0.022	2.133	
A110111	I1636	501-125	1 08 48.0	33 06 00	-2	5652	141	0	141	2.149	0.019	2.142	
U752	N420	501-127	1 09 23.6	31 51 28	-5	4980	173	0	173	2.238	0.018	2.231	
A110121	I1638	501-129	1 09 35.5	33 05 47	1	4782	168	44	166	2.220	0.020	2.213	
U776	N431	501-132	1 11 16.4	33 26 25	0	5713	157	27	156	2.193	0.021	2.186	
U792	I1652	502-009	1 12 09.7	31 41 03	-2	5067	113	0	113	2.053	0.020	2.046	
U804	N447	502-013	1 12 49.8	32 48 10	0	5586	152	31	150	2.176	0.020	2.169	
A110205	N468	502-029	1 17 00.0	32 30 07	1	5079	87	0	87	1.940	0.030	1.933	
U870	N472	502-034	1 17 40.4	32 26 55	-5	5280	261	70	255	2.407	0.014	2.400	
A110226	I1673	502-038	1 17 57.9	32 46 59	-5	5063	173	59	169	2.228	0.018	2.221	
U878	...	521-004	1 18 13.9	33 38 12	-2	4057	165	93	155	2.190	0.024	2.183	
A110232	...	502-043	1 18 16.9	33 07 02	-5	5208	126	126	126	2.160	0.018	2.093	
A110237	...	502-044	1 18 28.9	32 49 44	-5	5103	127	0	127	2.104	0.022	2.097	
A110246	I1679	502-048	1 18 55.5	33 14 07	0	5278	76	43	73	1.863	0.030	1.856	

TABLE 2. (continued)

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
U906	N483	502-050	1 19 07.0	33 15 41	0	4698	204	38	202	2.305	0.015	2.298	
A110268	...	521-011	1 19 59.4	33 42 27	0	5501	85	23	83	1.919	0.028	1.912	
A110272	...	...	1 20 02.7	33 15 23	-2	4278	133	92	121	2.083	0.026	2.076	
A110281	I1687	502-061	1 20 30.1	33 01 01	-5	5037	161	0	160	2.204	0.025	2.197	
U920	N495	502-058	1 20 07.1	33 12 40	1	4076	164	34	162	2.210	0.019	2.203	
U926	N499	502-059	1 20 22.8	33 12 01	-5	4358	258	0	258	2.412	0.016	2.405	1
U935	N504	502-064	1 20 39.1	32 56 38	-2	4242	192	60	188	2.274	0.020	2.267	
U938	N507	502-067	1 20 51.1	32 59 40	-5	4904	283	0	283	2.452	0.020	2.445	1
U939	N508	502-068	1 20 51.6	33 01 09	-5	5490	203	27	202	2.305	0.019	2.298	
A110833	I1689	502-070	1 20 58.9	32 47 43	-5	4540	129	0	129	2.111	0.022	2.104	
A110293	I1690	502-071	1 21 00.5	32 53 45	-2	4591	144	43	141	2.149	0.024	2.142	
A110296	...	502-072	1 21 09.3	33 03 10	-2	5009	139	0	139	2.143	0.021	2.136	
A110302	...	502-074	1 21 18.0	32 31 00	-2	5997	82	0	82	1.914	0.028	1.907	
U956	N515	502-077	1 21 48.8	33 12 45	-2	5101	225	88	215	2.332	0.016	2.325	
A110305	I1692	502-076	1 21 50.3	32 58 33	-2	5574	91	37	88	1.944	0.026	1.938	
U960	N517	502-079	1 21 54.2	33 10 08	-5	4150	186	40	184	2.265	0.015	2.258	
U988	N528	502-083	1 22 44.6	33 24 45	-2	4750	271	93	264	2.422	0.016	2.415	
U995	N529	521-023	1 22 50.0	34 27 14	-5	4792	244	63	240	2.380	0.017	2.373	1
A110325	...	502-084	1 23 22.9	33 08 44	-2	5059	98	21	97	1.987	0.022	1.980	
U1012	N531	521-024	1 23 28.4	34 29 40	1	4595	144	77	136	2.134	0.028	2.127	
A110327	N542	521-026	1 23 40.4	34 24 56	-2	4651	96	25	95	1.978	0.035	1.971	
U1058	N566	502-092	1 26 12.0	32 05 00	-2	5386	118	24	117	2.068	0.025	2.061	
A110536	...	522-008	1 46 07.2	34 59 09	0	4614	97	27	96	1.982	0.024	1.975	
U1269	...	522-009	1 46 11.0	34 44 05	-5	3819	130	0	130	2.114	0.026	2.106	
U1272	...	522-010	1 46 21.2	34 49 26	1	4967	138	0	138	2.140	0.014	2.132	
A110540	...	522-012	1 46 25.3	35 59 27	1	4069	81	37	79	1.898	0.028	1.890	
U1277	...	522-014	1 46 30.6	35 12 16	-2	4141	150	90	140	2.146	0.018	2.139	
U1283	N679	522-015	1 46 48.3	35 32 15	-2	5036	242	20	242	2.384	0.013	2.376	1
U1298	N687	522-017	1 47 37.7	36 07 25	-2	5072	232	24	231	2.364	0.014	2.356	1
U1308	...	522-022	1 47 55.3	36 01 45	-2	5212	223	0	223	2.348	0.022	2.341	
A110851	...	522-023	1 48 23.6	34 37 29	0	5022	95	0	95	1.978	0.022	1.970	
A110581	...	522-026	1 49 11.1	35 10 04	-2	4868	171	0	171	2.233	0.018	2.225	

TABLE 2. (continued)

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
U1336	N700	522-027	1 49 16.4	35 51 03	-2	4263	117	0	117	2.068	0.032	2.061	
A110586	...	522-030	1 49 20.6	35 47 24	-2	4579	142	0	142	2.152	0.022	2.145	
U1339	...	522-032	1 49 28.7	35 35 35	1	4081	153	24	152	2.182	0.019	2.174	
U1346	N703	522-037	1 49 43.2	35 55 28	-5	5555	269	116	257	2.410	0.019	2.402	
A111122	...	...	1 49 43.6	35 53 42	-5	5422	109	0	109	2.037	0.028	2.030	
U1345	N705	522-036	1 49 45.4	35 53 51	0	4497	172	48	169	2.228	0.017	2.220	
U1348	N708	522-039	1 49 49.9	35 54 22	-5	4837	212	33	210	2.322	0.033	2.315	1
A111123	...	...	1 49 54.3	35 54 20	-5	5492	63	...	63	1.799	0.041	1.792	
U1352	N712	522-043	1 50 11.9	36 34 32	-2	5339	252	142	238	2.377	0.020	2.369	
A110594	...	522-044	1 50 18.1	36 05 17	-2	4928	130	38	128	2.107	0.020	2.100	
U1353	...	522-046	1 50 25.8	36 42 35	-5	5210	220	107	210	2.322	0.015	2.315	
A110597	...	522-045	1 50 27.3	35 45 57	-5	4788	126	0	126	2.100	0.020	2.093	
U1358	N714	522-047	1 50 32.9	35 58 33	0	4403	264	87	255	2.407	0.016	2.399	
A110605	...	522-048	1 50 53.3	36 06 15	-5	4137	157	51	153	2.185	0.020	2.177	
U1363	N717	522-052	1 50 58.3	35 59 01	0	5099	138	70	129	2.111	0.024	2.103	
A110607	...	522-053	1 50 59.9	36 23 42	1	5679	106	42	101	2.004	0.026	1.997	
A110608	...	522-054	1 51 14.8	35 40 18	1	4292	96	56	90	1.954	0.029	1.947	
A110613	...	522-057	1 51 55.6	37 17 01	0	4553	109	0	109	2.037	0.022	2.030	
U1392	...	...	1 52 32.6	36 19 03	1	5248	87	0	87	1.940	0.028	1.932	
U1406	N732	522-076	1 53 30.1	36 33 32	0	5883	133	19	133	2.124	0.016	2.116	
U1415	...	522-080	1 53 46.5	36 08 17	0	4624	141	0	141	2.149	0.021	2.142	
A110683	...	522-085	1 54 39.3	37 12 43	-5	4519	99	0	99	1.996	0.025	1.988	
U1434	...	522-084	1 54 41.5	36 00 41	0	4566	113	0	113	2.053	0.026	2.046	
U1440	N759	522-087	1 54 53.1	36 06 00	-5	4646	240	0	240	2.380	0.017	2.373	1
A110690	...	522-091	1 55 06.9	37 20 06	-5	4808	186	75	179	2.253	0.016	2.245	
A110711	...	522-098	1 56 23.5	36 35 02	-5	4836	159	35	158	2.199	0.025	2.191	
U1475	I179	522-101	1 57 12.3	37 46 43	-5	4185	225	84	218	2.338	0.014	2.331	1
A180152	I2253	119-024	8 13 38.3	21 33 53	-2	4714	230	47	227	2.356	0.026	2.350	
A180170	N2553	119-031	8 14 40.0	21 03 32	-2	4695	199	92	192	2.283	0.028	2.277	
A180195	N2556	119-045	8 16 05.9	21 05 39	-5	4619	173	77	164	2.215	0.032	2.208	
U4330	N2557	119-048	8 16 15.2	21 35 34	-2	4833	183	43	181	2.258	0.032	2.251	
U4337	N2560	119-058	8 16 57.0	21 08 34	-2	4809	233	101	218	2.338	0.017	2.332	

TABLE 2. (continued)

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
U4347	N2563	119-065	8 17 40.7	21 13 40	-5	4545	271	72	268	2.428	0.015	2.422	1
A180225	N2569	119-067	8 18 26.4	21 01 37	-5	5124	161	64	158	2.199	0.024	2.192	
A180237	...	119-077	8 19 58.2	21 14 21	-2	4443	146	42	144	2.158	0.028	2.152	
U4384	I2341	119-081	8 20 46.3	21 35 49	-2	5141	142	28	141	2.149	0.030	2.143	
A180278	...	119-101	8 24 06.0	21 52 40	-5	4560	91	26	89	1.949	0.022	1.943	
A210505	...	97-021	11 33 51.1	20 05 42	-2	6648	127	0	127	2.104	0.030	2.103	
A210512	...	97-023	11 34 15.2	20 16 54	-5	6320	146	28	144	2.158	0.021	2.157	
U6642	N3805	127-024	11 38 05.9	20 37 13	-2	6534	216	80	212	2.326	0.032	2.325	
A210603	...	97-055	11 38 46.1	20 14 27	-5	6546	196	43	193	2.286	0.018	2.284	
U6663	N3821	127-032	11 39 33.0	20 35 38	0	5758	196	0	196	2.292	0.017	2.291	
A210639	...	97-074	11 40 23.3	20 21 42	-2	6036	176	79	169	2.228	0.020	2.227	
A210662	N3834	97-084	11 41 01.8	19 22 12	-2	6796	219	26	217	2.336	0.028	2.335	
A210666	...	127-040	11 41 07.4	20 32 47	-2	6895	166	54	162	2.210	0.019	2.208	
A210674	...	97-090B	11 41 21.9	20 14 00	-2	6148	179	59	175	2.243	0.020	2.242	
A210676	...	97-088	11 41 23.5	20 03 35	-2	5634	152	20	151	2.179	0.030	2.178	
A210679	N3841	97-096	11 41 24.0	20 14 27	-2	6313	143	41	141	2.149	0.032	2.148	
U6705	N3844	97-097	11 41 24.8	20 18 19	-5	6828	201	36	199	2.299	0.019	2.298	
U6704	N3842	97-095	11 41 26.4	20 13 40	-5	6263	304	22	304	2.483	0.032	2.482	1
A210683	N3845	97-100	11 41 29.9	20 16 35	-2	5653	179	30	177	2.248	0.032	2.247	
A210691	N3851	97-106	11 41 44.8	20 15 33	-2	6376	255	137	240	2.380	0.028	2.379	
A210708	...	97-112	11 41 54.4	20 21 13	-2	6713	138	19	137	2.137	0.017	2.136	
A211330	...	...	11 42 12.7	19 51 30	-5	6256	214	28	214	2.330	0.032	2.329	
A210729	...	97-124	11 42 20.7	20 00 11	-2	7771	163	0	163	2.212	0.018	2.211	
A210739	I2955	97-128	11 42 29.9	19 53 10	-2	6468	196	0	196	2.292	0.022	2.291	1
A210741	...	97-131	11 42 38.7	20 07 25	-2	7718	189	80	183	2.262	0.032	2.261	
A210749	...	127-048	11 42 51.8	21 05 40	-5	6973	288	57	285	2.455	0.028	2.454	
A210750	N3868	97-135	11 42 54.6	19 43 29	-2	6385	218	79	212	2.326	0.032	2.325	
U6731	N3867	97-134	11 42 55.4	19 41 14	0	7466	227	0	227	2.356	0.035	2.355	
A210765	...	97-143A	11 43 29.9	20 04 11	-5	5663	161	54	161	2.207	0.018	2.206	
A210765b	...	97-143B	11 43 29.9	20 04 11	-5	7170	84	...	78	1.892	0.030	1.891	
U6760	N3886	97-147	11 44 29.9	20 06 56	-5	5860	254	75	250	2.398	0.024	2.397	
A210795	...	97-155	11 45 28.0	20 17 10	-5	7237	139	62	135	2.130	0.026	2.129	

TABLE 2. (continued)

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
U6810	N3919	97-161	11 48 06.8	20 17 35	-5	6111	281	0	281	2.449	0.026	2.448	
U6851	N3937	127-088	11 50 07.9	20 54 38	-2	6720	308	0	308	2.489	0.024	2.487	
A210878	...	127-092	11 50 36.3	20 56 13	-5	6873	148	0	148	2.170	0.022	2.169	
U8038	N4798	160-013	12 52 29.1	27 41 11	-2	7851	169	0	169	2.228	0.033	2.228	
U8049	N4807	160-017	12 53 03.8	27 47 28	-5	6979	227	26	227	2.356	0.024	2.356	1,2
A221133	N4821	160-024	12 54 05.1	27 14 00	-2	7027	169	0	169	2.228	0.027	2.228	
U8065	N4827	160-028	12 54 18.0	27 26 55	-5	7610	290	0	289	2.461	0.024	2.461	2
U8070	N4839	160-039	12 54 58.3	27 46 11	-2	7318	245	0	245	2.389	0.024	2.389	1,2*
A221178	N4842	160-046	12 55 10.2	27 45 25	-5	7317	207	79	202	2.305	0.028	2.305	
U8086	N4849	160-056S	12 55 47.5	26 40 01	-5	5911	215	0	215	2.332	0.028	2.332	
A221216	...	160-062	12 55 53.6	29 23 53	-2	7867	140	0	138	2.140	0.021	2.140	
U8097	N4859	160-071	12 56 37.0	27 05 06	-2	7167	233	132	217	2.336	0.030	2.336	
A221266	I3957	160-217	12 56 42.8	28 02 21	-5	6376	149	22	148	2.170	0.033	2.170	1
U8103	N4874	160-231	12 57 11.1	28 13 53	-5	7189	204	0	204	2.310	0.024	2.310	1,2*
A221354	N4886	160-239	12 57 40.0	28 15 32	-5	6397	150	32	148	2.170	0.024	2.170	1,2
U8110	N4889	160-241	12 57 43.7	28 14 54	-5	6482	402	0	402	2.604	0.016	2.604	1,2*
A221410	I4041	160-254	12 58 16.4	28 16 06	-2	7090	116	0	116	2.064	0.024	2.064	3
A221422	...	160-091	12 58 28.2	28 38 00	-2	7647	204	43	202	2.305	0.026	2.305	
U8133	N4919	160-094	12 58 53.2	28 04 44	-2	7300	167	48	164	2.215	0.028	2.215	
U8135	N4922	160-096S	12 59 00.7	29 34 41	-5	7154	196	24	194	2.288	0.024	2.288	
U8154	N4931	160-118	13 00 37.0	28 18 07	-5	5418	213	94	205	2.312	0.022	2.312	
U8167	N4944	160-124	13 01 25.8	28 27 24	-2	6988	203	77	198	2.297	0.027	2.297	
U8175	N4952	160-129	13 02 35.4	29 23 28	-5	5963	228	0	228	2.358	0.024	2.358	2
A330131	...	...	23 11 47.5	9 39 54	0	17615	160	116	144	2.158	0.026	2.146	
A330144	...	406-031	23 12 16.3	7 26 44	1	4701	87	0	87	1.940	0.041	1.927	
U12454	...	406-033	23 12 37.7	9 24 27	-5	4833	159	33	158	2.199	0.026	2.186	
A330161	N7557	406-035	23 13 07.8	6 26 00	-2	3749	74	0	74	1.869	0.022	1.857	
U12464	N7562	406-039	23 13 25.1	6 24 53	-5	3605	243	0	243	2.386	0.017	2.373	1
U12509	N7611	406-066	23 17 04.9	7 47 24	-2	3303	201	44	199	2.299	0.015	2.286	
U12512	N7612	406-068	23 17 12.2	8 18 09	-2	3240	180	0	180	2.255	0.016	2.243	
A330238	...	406-069	23 17 25.0	8 17 12	1	3204	109	0	108	2.033	0.041	2.021	

TABLE 2. (continued)

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
A330246	N7617	406-072	23 17 37.0	7 53 31	-2	4156	114	37	111	2.045	0.022	2.033	1
U12523	N7619	406-073	23 17 42.6	7 55 57	-2	3765	340	20	339	2.530	0.019	2.518	1
U12526	N7623	406-075	23 17 58.0	8 07 20	-2	3680	181	0	181	2.258	0.020	2.245	
U12531	N7626	406-076	23 18 10.3	7 56 35	-5	3401	276	0	276	2.441	0.018	2.429	1
A330250	N7621	406-074	23 17 48.0	8 04 00	-5	3834	67	0	67	1.826	0.030	1.814	
A330257	...	...	23 18 24.0	7 54 45	-2	3470	78	0	78	1.892	0.024	1.880	
A330266	...	406-078	23 18 41.0	7 05 30	0	3155	169	38	168	2.225	0.018	2.213	
U12542	N7634	406-085	23 19 10.0	8 36 47	-5	3218	180	36	178	2.250	0.018	2.238	
U12575	N7648	406-096	23 21 22.2	9 23 37	1	3507	66	29	64	1.806	0.041	1.794	
A330499	...	476-052	23 27 34.8	25 44 14	-5	9462	177	0	177	2.248	0.020	2.252	
A331654	...	...	23 28 33.2	25 48 35	-5	9362	116	...	116	2.064	0.026	2.068	
A330553	...	476-071	23 32 08.9	26 26 49	-2	9687	192	20	192	2.283	0.024	2.287	
A331223	...	...	23 33 15.8	26 19 44	-2	9463	153	38	151	2.179	0.026	2.183	
A331231	...	...	23 33 56.0	27 07 17	-2	9146	168	52	165	2.217	0.022	2.221	
A331233	...	...	23 33 59.9	26 07 05	-5	9142	174	59	170	2.230	0.028	2.234	
A330595	...	476-079	23 34 49.9	26 36 04	-2	8501	202	43	199	2.299	0.024	2.303	
A330600	...	476-080	23 34 55.8	26 47 40	-2	9984	120	47	114	2.057	0.032	2.061	
A330602	...	...	23 34 57.5	26 21 21	-2	9683	136	38	134	2.127	0.026	2.131	
A330604	...	...	23 35 01.3	26 33 37	-5	8292	168	20	167	2.223	0.030	2.226	
A330614	...	476-081	23 35 14.7	26 13 41	-2	8593	163	63	159	2.201	0.022	2.205	
A330624	...	...	23 35 27.6	26 59 14	-5	10231	172	32	169	2.228	0.045	2.232	
A331239	...	...	23 35 42.1	26 13 02	-5	9206	180	108	166	2.220	0.022	2.224	
A330643	...	...	23 35 43.4	27 08 13	-2	9337	201	32	199	2.299	0.045	2.303	
A330647	...	476-084	23 35 45.7	26 32 25	-2	8727	180	49	178	2.250	0.020	2.254	
A330649	...	476-085	23 35 48.1	26 36 35	-5	9296	229	79	223	2.348	0.020	2.352	4
A330653	...	476-086	23 35 52.4	26 52 52	-5	9249	188	21	187	2.272	0.020	2.276	
A330658	I5341	476-087	23 35 56.5	26 42 29	-5	10779	191	0	191	2.281	0.020	2.285	4
A330660	...	476-090	23 35 58.8	26 42 06	-5	9503	228	0	228	2.358	0.021	2.362	4
U12716	N7720A	476-091S	23 35 59.1	26 45 14	-5	9031	277	0	277	2.442	0.018	2.446	1,4*
A331456	N7720B	476-091N	23 35 59.2	26 45 26	-5	8104	197	30	196	2.292	0.022	2.296	4*
A330667	...	476-092	23 36 04.1	26 42 09	-5	8493	242	49	240	2.380	0.021	2.384	4
A330668	...	476-090B	23 36 05.9	26 45 09	-2	8419	205	0	205	2.312	0.024	2.315	4

TABLE 2. (continued)

Name	NGC	CGCG	RA (1950)	Dec (1950)	T	V_{hel}	σ — (km s ⁻¹) —	V_{rot}	σ_c	$\log \sigma_c$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)
A330678	...	...	23 36 07.9	26 44 00	-5	9252	213	0	213	2.328	0.020	2.332	4
A330679	...	476-095B	23 36 11.3	26 56 17	-2	8444	104	...	104	2.017	0.019	2.021	
A330686	...	476-095A	23 36 13.3	26 56 17	-5	9236	237	25	236	2.373	0.037	2.377	
A330687	...	...	23 36 13.8	26 34 25	-2	9014	220	111	212	2.326	0.032	2.330	
A330696	...	476-096	23 36 20.3	26 59 26	-5	9269	262	32	261	2.417	0.016	2.420	
A330706	...	...	23 36 32.2	26 49 31	-5	9939	236	68	231	2.364	0.028	2.367	
A331248	...	...	23 36 41.9	27 16 25	-2	9267	203	115	193	2.286	0.026	2.289	
A330741	...	476-101	23 37 18.9	27 05 56	1	9921	188	31	187	2.272	0.022	2.276	
A330744	...	476-102	23 37 28.0	26 33 25	-5	8692	212	93	202	2.305	0.024	2.309	
U12727	N7728	476-103	23 37 30.3	26 51 22	-5	9397	338	41	338	2.529	0.019	2.533	
A330747	...	...	23 37 33.6	26 51 24	-2	9878	107	...	107	2.029	0.050	2.033	
A330754	...	476-105	23 37 48.5	27 17 03	-5	8668	233	77	229	2.360	0.024	2.364	
A331260	...	...	23 38 03.0	26 57 53	-5	10827	202	71	197	2.294	0.024	2.298	
U12733	...	476-107	23 38 16.1	26 33 29	-5	11704	233	0	233	2.367	0.024	2.371	
A330763	...	476-109	23 38 23.6	27 13 55	-5	9971	226	28	225	2.352	0.022	2.356	
U12744	N7735	476-115	23 39 42.0	25 57 00	-5	9594	260	0	260	2.415	0.020	2.419	
U12745	N7737	476-118	23 40 15.3	26 46 31	-2	7624	253	53	251	2.400	0.017	2.403	
A330785	...	476-120	23 40 25.2	27 11 40	-5	7616	234	0	234	2.369	0.019	2.373	
A331298	N7740	476-122	23 40 44.8	27 05 12	-5	7776	158	28	156	2.193	0.026	2.197	
A330798	...	476-123	23 41 01.1	27 02 04	-2	9126	275	75	272	2.435	0.020	2.438	
A330889	...	477-004	23 43 15.9	27 03 42	-2	9152	119	63	114	2.057	0.041	2.061	

References for Table 2.

(1) Faber et al. 1989; (2) Lucey et al. 1991b; (3) Dressler 1987; (4) Lucey et al. 1991a

Notes to Table 2.

U8070: σ also measured by Davies & Illingworth 1983, and Fisher, Illingworth & Franx 1995.U8103: σ also measured by Malumuth & Kirshner 1981, and Fisher, Illingworth & Franx 1995.U8110: σ also measured by Oegerle & Hoessel 1991, Tonry 1985, Tonry & Davis 1981, Davies & Illingworth 1983, Faber & Jackson 1976, and Fisher, Illingworth & Franx 1995.U12716: σ also measured by Tonry 1984, Malumuth & Kirshner 1981, Tonry & Davis 1981.331456: σ also measured by Tonry 1984.

TABLE 3. Spectroscopic parameters from the literature

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} – (km/s) –	σ	$\log \sigma$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
U683	N379	501-082	1 04 30.0	32 15 16	-2	5556	245	2.390	0.039	2.383	1
U682	N380	501-081	1 04 32.3	32 13 01	-5	4414	277	2.443	0.039	2.436	1
U688	N382	501-086	1 04 38.7	32 08 13	-5	5228	153	2.185	0.039	2.178	1
U689	N383	501-087	1 04 39.4	32 08 46	-2	5071	265	2.424	0.039	2.417	1
U687	N385	501-085	1 04 41.9	32 03 15	-5	5024	180	2.255	0.039	2.248	1
A110044	N386	501-088	1 04 45.5	32 05 43	-5	5540	61	1.785	0.039	1.778	1
U700	N392	501-094	1 05 37.1	32 52 00	-2	4672	261	2.416	0.039	2.409	1
U735	N410	501-118	1 08 11.6	32 53 10	-5	5294	321	2.507	0.039	2.500	1
U926	N499	502-059	1 20 22.8	33 12 01	-5	4375	237	2.374	0.039	2.367	1
A110284	N501	502-062	1 20 33.3	33 10 20	-5	4887	163	2.212	0.039	2.205	1
U938	N507	502-067	1 20 51.1	32 59 40	-5	4915	366	2.563	0.039	2.556	1
U995	N529	521-023	1 22 50.0	34 27 14	-5	4862	216	2.334	0.039	2.327	1
U1283	N679	522-015	1 46 48.3	35 32 15	-2	5054	232	2.365	0.039	2.357	1
U1298	N687	522-017	1 47 37.7	36 07 25	-2	5147	242	2.383	0.039	2.375	1
U1343N	N704A	522-034N	1 49 41.2	35 52 45	-5	4618	158	2.199	0.039	2.191	1
U1348	N708	522-039	1 49 49.9	35 54 22	-5	4827	242	2.383	0.039	2.375	1
U1440	N759	522-087	1 54 53.1	36 06 0	-5	4714	271	2.433	0.039	2.425	1
U1475	I179	522-101	1 57 12.3	37 46 43	-5	4209	214	2.330	0.039	2.322	1
U4347	N2563	119-065	8 17 40.7	21 13 40	-5	4642	261	2.416	0.039	2.409	1
U6701	N3837	97-089	11 41 21.0	20 10 21	-5	6248	258	2.412	0.039	2.411	1
A210679	N3841	97-096	11 41 24.0	20 14 27	-2	6363	210	2.322	0.039	2.321	1
U6704	N3842	97-095	11 41 26.4	20 13 40	-5	6237	282	2.451	0.039	2.450	1
U6723	N3862	97-127	11 42 29.1	19 53 05	-2	6462	262	2.418	0.039	2.417	1
A210739	I2955	97-128	11 42 29.9	19 53 10	-2	6345	188	2.274	0.039	2.273	1
U6735	N3873	97-137	11 43 10.2	20 03 08	-5	5438	243	2.386	0.039	2.385	1
U6852	N3940	127-089	11 50 11.8	21 16 06	-5	6506	208	2.318	0.039	2.317	1
A221044	...	159-083	12 47 16.2	27 10 00	-5	6887	210	2.322	0.025	2.322	2

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} – (km/s) –	σ	$\log \sigma$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
A221057	...	159-089	12 48 26.6	28 06 33	-5	7606	176	2.246	0.025	2.246	2
A221100	I832	159-105	12 51 30.0	26 43 18	-5	7094	217	2.336	0.025	2.336	2
U8028	N4789	160-008	12 51 53.1	27 20 25	-5	8369	272	2.435	0.025	2.435	2
U8049	N4807	160-017	12 53 03.8	27 47 28	-5	6941	208	2.318	0.039	2.318	1
						6973	212	2.326	0.025	2.326	2
A221121	I3900	160-019	12 53 15.6	27 31 37	-5	7115	280	2.447	0.025	2.447	2
U8057	N4816	160-021	12 53 46.8	28 00 56	-2	6931	210	2.322	0.025	2.322	2
A221129	I834	160-022	12 53 52.7	26 37 45	-5	6457	269	2.430	0.025	2.430	2
A221128	...	160-023	12 53 54.4	28 01 16	-5	6903	184	2.265	0.025	2.265	2
A221134	...	160-027	12 54 01.3	28 06 03	-5	6275	178	2.250	0.025	2.250	2
A221132	...	D140	12 54 04.3	28 12 47	-5	6689	177	2.248	0.025	2.248	2
A221135	N4824	...	12 54 08.3	27 48 40	-5	7121	164	2.215	0.025	2.215	2
U8065	N4827	160-028	12 54 18.0	27 26 55	-5	7599	279	2.446	0.025	2.446	2
A221151	...	160-033	12 54 25.5	27 10 07	-2	6273	85	1.929	0.025	1.929	2
A221164	...	160-037	12 54 45.4	27 44 20	-5	7480	237	2.375	0.025	2.375	2
U8070	N4839	160-039	12 54 58.3	27 46 11	-2	7446	259	2.413	0.039	2.413	1
						7378	298	2.474	0.025	2.474	2
U8072	N4841A	160-044S	12 55 07.4	28 44 50	-5	6784	264	2.422	0.039	2.422	1
A221175	N4840	160-042	12 55 07.5	27 52 55	-5	6097	250	2.398	0.025	2.398	2
U8073	N4841B	160-044N	12 55 09.5	28 45 21	-5	6793	229	2.360	0.039	2.360	1
A221188	...	160-049	12 55 22.9	28 27 12	-5	7245	179	2.253	0.025	2.253	2
A221193	...	D238	12 55 28.7	28 46 21	-5	7339	110	2.053	0.025	2.053	2
A221220	N4850	160-063	12 55 57.1	28 14 25	-5	6044	180	2.255	0.025	2.255	2
A221226	...	160-065	12 56 05.4	28 17 13	-2	7188	188	2.274	0.025	2.274	2
A221242	N4854	160-070	12 56 22.4	27 56 45	-2	8077	175	2.243	0.039	2.243	1
						8394	204	2.310	0.025	2.310	2
A221247	...	D181	12 56 26.3	28 21 21	-2	5932	142	2.152	0.039	2.152	3
A221248	I3947	160-211	12 56 27.2	28 03 33	-2	5677	148	2.170	0.025	2.170	2
A221252	...	D136	12 56 30.6	28 14 12	-2	5668	183	2.262	0.039	2.262	1
						5706	137	2.137	0.025	2.137	2
A221256	...	D135	12 56 35.7	28 14 23	-5	8320	85	1.929	0.025	1.929	2
A221259	N4860	160-215	12 56 39.3	28 23 47	-5	7966	265	2.423	0.025	2.423	2
A221261	...	D180	12 56 40.3	28 20 05	-5	6396	116	2.064	0.039	2.064	3

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} – (km/s) –	σ	$\log \sigma$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
A221263	I3955	160-216	12 56 41.1	28 16 06	-2	7895	181	2.258	0.039	2.258	3
A221266	I3957	160-217	12 56 42.8	28 02 21	-5	6309	150	2.176	0.039	2.176	1
A221267	I3960	160-219	12 56 43.1	28 07 35	-5	6709	175	2.243	0.039	2.243	3
A221268	I3959	160-218	12 56 43.5	28 03 16	-5	7084	195	2.290	0.025	2.290	2
						7121	200	2.301	0.039	2.301	1
A221274	N4864	160-221	12 56 48.4	28 14 54	-5	6760	198	2.297	0.039	2.297	1
A221275	I3963	160-220	12 56 48.6	28 02 43	-5	6675	129	2.111	0.039	2.111	3
A221277	N4867	160-222	12 56 50.6	28 14 33	-5	4818	222	2.346	0.039	2.346	1
U8100	N4865	160-224	12 56 54.9	28 21 14	-5	4643	234	2.369	0.039	2.369	3
A221284	...	D106	12 56 58.1	28 10 05	-5	5092	159	2.201	0.039	2.201	3
A221285	N4869	160-225	12 56 58.2	28 10 54	-5	6703	205	2.312	0.039	2.312	1
A221289	...	D67	12 57 00.2	28 00 46	-5	6008	148	2.170	0.039	2.170	3
A221290	...	D132	12 57 00.8	28 14 23	-5	7663	127	2.104	0.039	2.104	3
A221291	...	D157	12 57 00.9	28 14 42	-5	6083	126	2.100	0.039	2.100	3
A221293	...	D156	12 57 01.8	28 16 15	-2	6724	107	2.029	0.025	2.029	2
A221295	...	D177	12 57 04.1	28 18 43	-2	6871	99	1.996	0.039	1.996	3
A221296	I3967	160-226	12 57 04.7	28 07 15	-2	6950	251	2.400	0.039	2.400	3
A221298	N4871	160-227	12 57 05.5	28 13 42	-2	7113	168	2.225	0.039	2.225	3
A221301	I3973	160-228	12 57 06.3	28 09 20	-2	4745	208	2.318	0.039	2.318	3
A221302	...	D176	12 57 06.9	28 19 05	-2	5579	161	2.207	0.039	2.207	3
A221303	N4873	160-229	12 57 08.1	28 15 19	-2	5662	150	2.176	0.039	2.176	3
A221304	N4872	160-230	12 57 09.6	28 13 09	-5	7145	212	2.326	0.039	2.326	1
U8103	N4874	160-231	12 57 11.1	28 13 53	-5	7176	245	2.389	0.039	2.389	1
						7200	279	2.446	0.025	2.446	2
A221313	N4875	160-232	12 57 13.2	28 10 43	-5	7897	182	2.260	0.039	2.260	3
A221317	...	D128	12 57 15.1	28 13 32	-2	7681	104	2.017	0.039	2.017	3
A221323	...	160-233	12 57 17.9	28 11 47	-5	6948	183	2.262	0.025	2.262	2
						6804	167	2.223	0.039	2.223	1
A221329	...	D153	12 57 19.2	28 15 57	-5	6640	135	2.130	0.039	2.130	1
A221331	N4876	160-234	12 57 19.8	28 11 02	-2	6741	194	2.290	0.025	2.290	2
						6629	182	2.260	0.039	2.260	1
A221332	...	160-235	12 57 21.5	28 07 42	-5	8028	126	2.100	0.039	2.100	3
A221334	...	D152	12 57 22.5	28 14 46	-2	9371	156	2.193	0.039	2.193	3

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} – (km/s) –	σ	$\log \sigma$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
A221335	...	160-079	12 57 23.0	27 58 51	-2	8219	161	2.207	0.039	2.207	3
A221343	...	D193	12 57 30.8	28 24 00	-2	7544	120	2.079	0.039	2.079	1
A221345	N4883	160-237	12 57 31.5	28 18 24	-2	7961	171	2.233	0.039	2.233	3
U8106	N4881	160-238	12 57 32.8	28 31 08	-5	6730	210	2.322	0.025	2.322	2
A221354	N4886	160-239	12 57 40.0	28 15 32	-5	6218	164	2.215	0.039	2.215	1
						6421	133	2.124	0.025	2.124	2
A221357	...	D65	12 57 41.6	28 02 46	-5	6053	115	2.061	0.039	2.061	3
A221362	I4011	160-242	12 57 42.0	28 16 33	-2	7142	106	2.025	0.039	2.025	1
A221364	I4012	160-244	12 57 43.4	28 20 59	-2	7218	179	2.253	0.039	2.253	1
U8110	N4889	160-241	12 57 43.7	28 14 54	-5	6497	381	2.581	0.039	2.581	1
						6515	414	2.617	0.025	2.617	2
A221365	...	D207	12 57 43.8	28 26 31	-2	6764	147	2.167	0.039	2.167	1
A221372	...	D173	12 57 48.4	28 20 48	-5	7384	137	2.137	0.039	2.137	3
A221376	I4021	160-246	12 57 50.3	28 18 45	-5	5789	158	2.199	0.039	2.199	1
A221377	N4894	160-247	12 57 52.1	28 14 20	-2	4587	91	1.959	0.039	1.959	3
A221380	N4898A	160-248W	12 57 53.1	28 13 35	-2	6848	201	2.303	0.039	2.303	1
U8113	N4895	160-249	12 57 53.5	28 28 15	-5	8406	213	2.328	0.039	2.328	3
A221382	N4898B	160-248E	12 57 53.8	28 13 41	-2	6349	140	2.146	0.039	2.146	1
A221386	I4026	160-250	12 57 57.7	28 19 04	-5	8198	140	2.146	0.039	2.146	3
A221390	...	D27	12 58 01.6	27 47 11	-5	7825	106	2.025	0.025	2.025	2
A221392	...	D119	12 58 03.5	28 13 38	-2	6927	153	2.186	0.039	2.186	3
A221403	...	160-251	12 58 11.1	28 25 02	-2	5526	91	1.959	0.039	1.959	3
A221404	...	D146	12 58 14.2	28 17 11	-2	7020	105	2.021	0.039	2.021	3
A221408	N4906	160-253	12 58 15.2	28 11 42	-5	7505	168	2.225	0.039	2.225	1
A221410	I4041	160-254	12 58 16.4	28 16 06	-2	7087	133	2.124	0.039	2.124	3
A221412	I4042	160-255	12 58 18.3	28 14 32	-2	6255	162	2.210	0.039	2.210	3
A221413	...	D116	12 58 18.5	28 14 04	-5	8366	130	2.114	0.039	2.114	3
A221416	...	D191	12 58 20.4	28 22 08	-2	6611	89	1.949	0.039	1.949	3
A221419	I4045	160-256	12 58 24.5	28 21 40	-2	6855	211	2.324	0.039	2.324	1
A221420	N4908	160-258	12 58 27.2	28 18 48	-5	8742	206	2.314	0.025	2.314	2
U8129	I4051	160-259	12 58 30.1	28 16 42	-5	4964	223	2.348	0.039	2.348	1
A221429	...	160-261	12 58 35.0	28 10 13	-2	6907	147	2.167	0.039	2.167	3
A221434	...	160-092	12 58 44.7	28 05 16	-2	5975	158	2.199	0.025	2.199	2

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} – (km/s) –	σ	$\log \sigma$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
A221438	...	D204	12 58 57.8	28 27 59	-2	7667	135	2.130	0.025	2.130	2
A221442	N4923	160-097	12 59 07.4	28 07 02	-2	5525	209	2.320	0.025	2.320	2
U8137	I843	160-099	12 59 09.5	29 23 58	-5	7387	255	2.407	0.025	2.407	2
A221453	...	160-100	12 59 26.2	28 09 47	-2	7604	192	2.283	0.025	2.283	2
U8142	N4926	160-103	12 59 29.7	27 53 35	-2	7874	266	2.425	0.025	2.425	2
A221455	N4927	160-105	12 59 33.6	28 16 35	-5	7747	292	2.465	0.025	2.465	2
A230015	N4929	160-113	13 00 20.5	28 18 54	-2	6234	191	2.281	0.025	2.281	2
A230042	I4133	160-123	13 01 26.6	28 15 23	-2	6377	178	2.250	0.025	2.250	2
U8175	N4952	160-129	13 02 35.4	29 23 28	-5	5948	191	2.281	0.025	2.281	2
U8178	N4957	160-130	13 02 48.4	27 50 15	-5	6922	249	2.396	0.025	2.396	2
A230086	N4971	160-140	13 04 30.9	28 48 46	-5	6292	184	2.265	0.025	2.265	2
U8260	N5004	160-157	13 08 39.3	29 54 08	-5	7061	244	2.387	0.025	2.387	2
U12464	N7562	406-039	23 13 25.1	6 24 53	-5	3608	243	2.385	0.039	2.373	1
A330246	N7617	406-072	23 17 37.0	7 53 31	-2	4072	139	2.142	0.039	2.130	1
U12523	N7619	406-073	23 17 42.6	7 55 57	-2	3747	337	2.528	0.039	2.516	1
U12531	N7626	406-076	23 18 10.3	7 56 35	-5	3450	234	2.369	0.039	2.357	1
A330595	...	476-079	23 34 49.9	26 36 04	-2	8525	285	2.455	0.040	2.459	4
A330600	...	476-080	23 34 55.8	26 47 40	-2	10005	123	2.090	0.040	2.094	4
A330615	...	...	23 35 16.4	26 42 14	-5	11056	118	2.072	0.040	2.076	4
A330621	...	...	23 35 25.1	26 49 24	-5	10173	105	2.021	0.040	2.025	4
A330624	...	...	23 35 27.6	26 59 14	-5	10264	179	2.253	0.040	2.257	4
A331239	...	...	23 35 42.1	26 13 02	-5	9181	160	2.204	0.040	2.208	4
A330643	...	...	23 35 43.4	27 08 13	-2	9324	248	2.394	0.040	2.398	4
A330648	...	...	23 35 47.5	26 45 33	-5	9538	174	2.241	0.040	2.244	4
A330649	...	476-085	23 35 48.1	26 36 35	-5	9293	239	2.378	0.040	2.382	4
A330651	...	...	23 35 50.3	26 46 26	-5	10137	142	2.152	0.040	2.156	4
A330653	...	476-086	23 35 52.4	26 52 52	-5	9109	154	2.188	0.040	2.191	4
A330658	I5341	476-087	23 35 56.5	26 42 29	-5	10765	234	2.369	0.040	2.373	4
A330662	...	476-089	23 35 58.2	26 32 29	-5	9485	154	2.188	0.040	2.191	4
A330660	...	476-090	23 35 58.8	26 42 06	-5	9514	209	2.320	0.040	2.324	4
U12716	N7720A	476-091S	23 35 59.1	26 45 14	-5	9141	305	2.484	0.039	2.488	1

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} $-(km/s) -$	σ	$\log \sigma$	ϵ_σ	$\log \sigma_n$	ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
A331456	N7720B	476-091N	23 35 59.2	26 45 26	-5	9083	353	2.548	0.040	2.551	4
A330665	...	...	23 36 02.9	26 45 28	-2	8117	158	2.199	0.040	2.202	4
A330667	...	476-092	23 36 04.1	26 42 09	-5	9985	184	2.265	0.040	2.269	4
A330668	...	476-090B	23 36 05.9	26 45 09	-2	8515	215	2.332	0.040	2.336	4
A330678	...	...	23 36 07.9	26 44 00	-5	8470	215	2.332	0.040	2.336	4
A330680	...	...	23 36 11.7	26 44 00	-5	9288	221	2.344	0.040	2.348	4
A330680	...	...	23 36 11.7	26 50 29	-5	9524	175	2.243	0.040	2.247	4
A330687	...	...	23 36 13.8	26 34 25	-2	9035	170	2.230	0.040	2.234	4
A330688	...	...	23 36 15.9	26 53 43	-2	9790	142	2.152	0.040	2.156	4
A330691	...	...	23 36 18.9	26 50 45	-2	8966	188	2.274	0.040	2.278	4
A330696	...	476-096	23 36 20.3	26 59 26	-5	9293	273	2.436	0.040	2.440	4
A330699	...	...	23 36 23.5	26 46 08	-5	9654	135	2.130	0.040	2.134	4
A330700	...	...	23 36 25.8	26 53 04	-5	9194	151	2.179	0.040	2.183	4
A330706	...	...	23 36 32.2	26 49 31	-5	9981	205	2.312	0.040	2.315	4
A330741	...	476-101	23 37 18.9	27 05 56	1	9930	171	2.233	0.040	2.237	4
A330744	...	476-102	23 37 28.0	26 33 25	-5	8685	242	2.384	0.040	2.388	4
U12727	N7728	476-103	23 37 30.3	26 51 22	-5	9408	314	2.497	0.040	2.501	4
A330746	...	...	23 37 32.6	26 53 23	-5	8990	165	2.217	0.040	2.221	4
A330754	...	476-105	23 37 48.5	27 17 03	-5	8706	257	2.410	0.040	2.414	4
A331260	...	...	23 38 03.0	26 57 53	-5	10865	254	2.405	0.040	2.409	4
U12733	...	476-107	23 38 16.1	26 33 29	-5	11704	256	2.408	0.040	2.412	4
A331265	...	...	23 38 20.0	26 33 13	-5	8738	161	2.207	0.040	2.211	4
U12744	N7735	476-115	23 39 42.0	25 57 00	-5	9580	277	2.442	0.040	2.446	4

References for Table 3.

(1) Faber et al. 1989; (2) Lucey et al. 1991b; (3) Dressler 1987; (4) Lucey et al. 1997